



HAL
open science

Double-scale theory

Michel Gondran, Alexandre Gondran

► **To cite this version:**

| Michel Gondran, Alexandre Gondran. Double-scale theory. 2023. hal-03960495

HAL Id: hal-03960495

<https://hal-enac.archives-ouvertes.fr/hal-03960495>

Preprint submitted on 27 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Double-scale theory

Michel Gondran

Académie Européenne Interdisciplinaire des Sciences, Paris, France
michel.gondran@polytechnique.org

Alexandre Gondran

École Nationale de l'Aviation Civile, Toulouse, France
alexandre.gondran@recherche.enac.fr

January 2023

Abstract

We present a new interpretation of quantum mechanics, called the double-scale theory, which expands on the de Broglie-Bohm (dBB) theory.

It is based, for any quantum system, on the simultaneous existence of two wave functions in the laboratory reference frame : an external wave function and an internal one. The external wave function is the wave function of the center-of-mass, as for the dBB theory : the wave is a field that pilots the center-of-mass of the quantum system. The external wave spreads out in space over time. Mathematically, the Schrödinger equation converges to the Hamilton-Jacobi statistical equations when the Planck constant tends towards zero and the Newton trajectories are therefore approximations of the dBB trajectories.

The simultaneous existence of an internal wave function, in addition to the external one, is the original element of our theory. This internal wave corresponds to the interpretation proposed by Edwin Schrödinger for whom the particle is extended. Then, the internal wave remains confined in space. It converges, when $\hbar \rightarrow 0$, to a Dirac distribution. Moreover, the configuration space of dimension $3N$ of the internal wave function can be written as the product of N individual internal wave functions of dimension 3.

1 Introduction

The theory of the double solution is an interpretation of quantum mechanics, which Louis de Broglie outlined as early as 1926 [1], and which he continued to work on for the last 30 years of his life [2, 3]. These studies remained for a long time a research program [4]. The pilot wave that he presented at the Solvay Congress in 1927 [5] and that was again found by Bohm in 1952 [6] and subsequently developed by the latter and his students [7, 8], was viewed by de Broglie as merely a temporary solution. He believed that there were two solutions to the Schrödinger equation. One is the “classical” wave function and corresponds to the pilot wave theory. The second, which he never found, should be a “Dirac function” which corresponds to a singularity in the wave function and which represents the “real” particle. De Broglie considered these two wave functions to be intimately related.

Following de Broglie’s idea, we presented recently [9] a new interpretation of quantum mechanics that defined two waves functions for any quantum system : an external one and an internal one. In this paper, we clarify this interpretation, which we name the double-scale theory, and in particular we demonstrate that the two wave functions exist simultaneously in the same reference frame, that of the laboratory.

These two functions have very different physical behaviors : the external wave function spreads out over time contrary to the internal wave function which remains confined. To justify the distinction

between these two wave functions, we demonstrate that the mathematical convergences are fundamentally different when $\hbar \rightarrow 0$. For the external wave function, its phase and the square of its modulus converge towards an action and a classical density satisfying the Hamilton-Jacobi statistical equations. For the internal wave function, the square of its modulus converges to a Dirac distribution, thus losing the internal structure of the atom or molecule.

We deduce very different interpretations for these two wave functions. By mathematical continuity, we show that the phase of the external wave “drives” the center-of-mass of the quantum system and corresponds to the de Broglie-Bohm interpretation restricted to the external wave function. The most likely interpretation of the internal wave is the one proposed by Schrödinger for which the particles are extended.

The plan of the paper is as follows. In section 2, we recall the well-known existence of the two wave functions: the center-of-mass wave function in the laboratory reference frame and the relative wave function in the center-of-mass reference frame.

In section 3, we show that the center-of-mass wave function in the laboratory reference frame is an external wave function that corresponds to a non-local field that “pilots” the center-of-mass of the corpuscle and that converges to classical mechanics when we make Planck’s constant tends to zero. This is the de Broglie-Bohm interpretation restricted to the external wave function.

In section 4, from the value of the center-of-mass and of the relative wave function, we define an internal wave function in the laboratory reference frame, that is local and corresponds to the corpuscle in the field-corpuscle duality. This is the Schrödinger interpretation restricted to the internal wave function. Moreover, the $3N$ dimensional configuration space of the internal wave function can be written as the product of N individual internal wave functions in the 3 dimensional space.

In section 5, the simultaneous existence of these two functions allows us to understand and explain simply many phenomena of quantum mechanics, such as wave-particle duality, quantum measurements and the nonlocality of the EPR-B experiment.

2 Center-of-mass wave and relative wave

Since the beginning of quantum mechanics, two types of variables have been distinguished in order to study the dynamics of atoms and molecules: internal variables and external variables. Roughly, the external variables correspond to the external dynamics of the atom, i.e. the motion of its center-of-mass and the orientation of the reference frame that is linked to it. The internal variables describe the structure of the atom or molecule in the reference frame of its center-of-mass.

These external and internal variables are linked by the well known decomposition of the total wave function into the center-of-mass wave function in the laboratory reference frame and the relative wave function in the center-of-mass reference frame. By rewriting the relative wave function in the laboratory reference frame, we obtain the so-called internal wave function. The double-scale theory is built on the existence of these two wave functions (external and internal) defined in the laboratory reference frame and by their interpretation. In this section we recall this decomposition for a quantum system with N bodies (atoms or molecules) where the decomposition is exact.

Let us consider a system of N spinless particles of masses m_j and charges q_j (with $j = 1..N$), of coordinates \mathbf{x}_j , subjected to an external gravitational field $V_g(\mathbf{x}_j)$ and a zero external electric field, and to mutual interactions described by the potentials $U_{jk}(|\mathbf{x}_j - \mathbf{x}_k|), \forall j, k = 1..N, j \neq k$. This quantum system is then described by the wave function $\Psi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N, t)$ which verifies the Schrödinger equation:

$$i\hbar \frac{\partial \Psi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N, t)}{\partial t} = H \Psi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N, t) \quad (1)$$

with the Hamiltonian:

$$H = \sum_{j=1}^N \left(\frac{\mathbf{p}_j^2}{2m_j} + m_j V_g(\mathbf{x}_j) \right) + \sum_{j=1}^N \sum_{\substack{k=1 \\ k \neq j}}^N U_{jk}(|\mathbf{x}_j - \mathbf{x}_k|) \quad (2)$$

and the initial condition :

$$\Psi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N, 0) = \Psi_0(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N). \quad (3)$$

We separate the motion of these N particles from the motion of their center-of-mass: let $\mathbf{x}_G = (\sum_{j=1}^N m_j \mathbf{x}_j) / (\sum_{j=1}^N m_j)$ be the position of the center-of-mass, $\mathbf{x}'_j = \mathbf{x}_j - \mathbf{x}_G$ be the relative position of the j particle and $M = \sum_{j=1}^N m_j$ be the total mass.

Then the Hamiltonian H is written as a function of the total impulse ($\mathbf{p}_G = \sum_{i=1}^N \mathbf{p}_i$) and relative impulses ($\mathbf{p}'_i = \mathbf{p}_i - m_i/M \mathbf{p}_G$) taking into account small variations of the gravitational field $V_g(\mathbf{x}_j) \simeq V_g(\mathbf{x}_G) + \mathbf{x}'_j \nabla V_g(\mathbf{x}_G)$:

$$H = \left(\frac{\mathbf{p}_G^2}{2M} + M V_g(\mathbf{x}_G) \right) + \sum_{i=1}^N \left(\frac{\mathbf{p}'_i{}^2}{2m_i} + \sum_{\substack{j=1 \\ j \neq i}}^N U_{ij}(|\mathbf{x}'_i - \mathbf{x}'_j|) \right) = H^{ext} + \sum_{i=1}^N H_i^{int}. \quad (4)$$

We note that the interaction of the gravitational field related to its local variation does not intervene because $\sum_{i=1}^N m_i (\mathbf{x}_i - \mathbf{x}_G) = 0$.

Proposition 1 - *If the initial wave function $\Psi_0(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N)$ factors to the form:*

$$\Psi_0(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N) = \psi_0(\mathbf{x}_G) \varphi_0(\mathbf{x}'_1, \mathbf{x}'_2, \dots, \mathbf{x}'_N). \quad (5)$$

then $\Psi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N, t)$, a solution to (1),(2) and (3), is written as the product of the center-of-mass wave function $\psi(\mathbf{x}_G, t)$ and the relative function $\varphi(\mathbf{x}'_1, \mathbf{x}'_2, \dots, \mathbf{x}'_N, t)$:

$$\Psi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N, t) = \psi(\mathbf{x}_G, t) \varphi(\mathbf{x}'_1, \mathbf{x}'_2, \dots, \mathbf{x}'_N, t) \quad (6)$$

where $\psi(\mathbf{x}_G, t)$ is the solution to the Schrödinger equations of the center-of-mass:

$$i\hbar \frac{\partial \psi(\mathbf{x}_G, t)}{\partial t} = -\frac{\hbar^2}{2M} \Delta_{\mathbf{x}_G} \psi(\mathbf{x}_G, t) + M V_g(\mathbf{x}_G) \psi(\mathbf{x}_G, t) \quad (7)$$

with the initial condition:

$$\psi(\mathbf{x}_G, 0) = \psi_0(\mathbf{x}_G) \quad (8)$$

and where $\varphi(\mathbf{x}'_1, \mathbf{x}'_2, \dots, \mathbf{x}'_N, t)$ is the Schrödinger relative equations:

$$i\hbar \frac{\partial \varphi(\mathbf{x}'_1, \mathbf{x}'_2, \dots, \mathbf{x}'_N, t)}{\partial t} = -\sum_{i=1}^N \left(\frac{\hbar^2}{2m_i} \Delta_{\mathbf{x}'_i} + \sum_{\substack{j=1 \\ j \neq i}}^N U_{ij}(|\mathbf{x}'_i - \mathbf{x}'_j|) \right) \varphi(\mathbf{x}'_1, \mathbf{x}'_2, \dots, \mathbf{x}'_N, t) \quad (9)$$

with the initial condition:

$$\varphi(\mathbf{x}'_1, \mathbf{x}'_2, \dots, \mathbf{x}'_N, 0) = \varphi_0(\mathbf{x}'_1, \mathbf{x}'_2, \dots, \mathbf{x}'_N). \quad (10)$$

We have considered the case of a decomposition of the total wave function as the product of the center-of-mass wave function in the laboratory reference frame and the relative wave function in the center-of-mass reference frame. This example highlights the simultaneous existence of these two functions. We postulate that the decomposition of the total wave function of an N -body quantum system $\Psi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N, t)$ into two waves functions $\psi(\mathbf{x}_G, t)$ and $\varphi(\mathbf{x}'_1, \mathbf{x}'_2, \dots, \mathbf{x}'_N, t)$ is not only a mathematical tool to simplify the calculations, but that these two waves physically exist at all times and simultaneously.

When the quantum system is not composed of several particles but corresponds to a single particle, like a free electron, we postulate that these two wave functions exist simultaneously, even if the mathematical decomposition is no longer necessary.

Hypothesis 1 - Hypothesis for a single particle: A single particle, like a free electron is also simultaneously described by two wave functions: an external wave function $\psi(\mathbf{x}_G, t)$ related to its center-of-mass and a relative wave function $\varphi(\mathbf{x}', t)$. This hypothesis assumes that the particle is not point-like and has an extension represented by the relative wave function; this is the Schrödinger conjecture of a single particle, which we develop in section 4.

The center-of-mass wave function and the relative wave function are defined in different reference frames. We will call the center-of-mass wave function the external wave function and the relative wave function the internal wave function when it is rewritten (see section 4) to be considered in the laboratory reference frame as the external wave function. These two wave functions, external and internal, have very different behaviors: the external wave function spreads over time and is non-local, while the internal wave function remains confined and is local. They will give rise to very different interpretations as we will see in the next two sections.

3 The external wave function: the field of the field-corpuscule duality

To interpret the external wave function, let us study its convergence to classical mechanics when we make Planck's constant tends towards zero.

3.1 Convergence of the external wave function

Let us consider an external wave function verifying the external Schrödinger equations (7) and (8) and make the semi-classical change of variable $\psi(\mathbf{x}_G, t) = \sqrt{\rho^{\hbar}(\mathbf{x}_G, t)} \exp\left(i \frac{S^{\hbar}(\mathbf{x}_G, t)}{\hbar}\right)$. The density $\rho^{\hbar}(\mathbf{x}_G, t)$ and the action $S^{\hbar}(\mathbf{x}_G, t)$ then verify the Madelung equations [10] (1926):

$$\frac{\partial S^{\hbar}(\mathbf{x}_G, t)}{\partial t} + \frac{(\nabla S^{\hbar}(\mathbf{x}_G, t))^2}{2M} + V(\mathbf{x}_G) - \frac{\hbar^2}{2M} \frac{\Delta \sqrt{\rho^{\hbar}(\mathbf{x}_G, t)}}{\sqrt{\rho^{\hbar}(\mathbf{x}_G, t)}} = 0 \quad \forall (\mathbf{x}_G, t) \in \mathbb{R}^3 \times \mathbb{R}^+ \quad (11)$$

$$\frac{\partial \rho^{\hbar}(\mathbf{x}_G, t)}{\partial t} + \text{div} \left(\rho^{\hbar}(\mathbf{x}_G, t) \frac{\nabla S^{\hbar}(\mathbf{x}_G, t)}{m} \right) = 0 \quad \forall (\mathbf{x}_G, t) \in \mathbb{R}^3 \times \mathbb{R}^+ \quad (12)$$

with the initial conditions

$$\rho^{\hbar}(\mathbf{x}_G, 0) = \rho_0^{\hbar}(\mathbf{x}_G) \quad \text{and} \quad S^{\hbar}(\mathbf{x}_G, 0) = S_0^{\hbar}(\mathbf{x}_G) \quad \forall \mathbf{x}_G \in \mathbb{R}^3 \quad (13)$$

Here, $V(\mathbf{x}_G) = MV_g(\mathbf{x}_G)$ is the potential of (7). Let us now study the convergence of the density $\rho^{\hbar}(\mathbf{x}_G, t)$ and the action $S^{\hbar}(\mathbf{x}_G, t)$ of the Madelung equations, when the Planck constant \hbar is made to tend to 0. We will restrict ourselves to “**prepared non-discerned quantum systems**”.

Definition 1 - A quantum system, subjected to a potential $V(\mathbf{x})$, is said to be a **prepared non-discerned quantum systems**, if we know at the initial time, only the density of the initial probability $\rho_0^{\hbar}(\mathbf{x}_G)$ and the initial action $S_0^{\hbar}(\mathbf{x}_G)$ of its external wave function, and that these are functions ρ_0 and $S_0(\mathbf{x}_G)$ which are independent of \hbar .

It is the case of a set of particles without interaction between each other and prepared in the same way: sources of free particles or in a linear field as in the Shimizu [11] experiment with cold atoms, sources of fullerenes, neutrons, electrons or C_{60} in a Young's slit experiment. One can then demonstrate [9] theorem below:

THEOREM 1 [9] - When \hbar tends to 0, if the external wave function is prepared non-discerned, the density $\rho^{\hbar}(\mathbf{x}_G, t)$ and the action $S^{\hbar}(\mathbf{x}_G, t)$, solutions to Madelung equations (11-13), converge to $\rho(\mathbf{x}_G, t)$ and $S(\mathbf{x}_G, t)$, solutions to Hamilton-Jacobi statistical equations:

$$\frac{\partial S(\mathbf{x}_G, t)}{\partial t} + \frac{(\nabla S(\mathbf{x}_G, t))^2}{2m} + V(\mathbf{x}_G) = 0 \quad (14)$$

$$S(\mathbf{x}_G, 0) = S_0(\mathbf{x}_G) \quad (15)$$

$$\frac{\partial \rho(\mathbf{x}_G, t)}{\partial t} + \text{div} \left(\rho(\mathbf{x}_G, t) \frac{\nabla S(\mathbf{x}_G, t)}{m} \right) = 0 \quad (16)$$

$$\rho(\mathbf{x}_G, 0) = \rho_0(\mathbf{x}_G) \quad (17)$$

3.2 Action-particle duality in classical mechanics

This Hamilton-Jacobi action $S(\mathbf{x}, t)$ is written from the Hamilton-Jacobi action at the initial time $S_0(\mathbf{x})$ and from the Euler-Lagrange actions, $S_{EL}(\mathbf{x}_0; \mathbf{x}, t)$, between all the possible paths from $(\mathbf{x}_0, 0)$ to (\mathbf{x}, t)

$$S(\mathbf{x}, t) = \min_{\mathbf{x}_0} (S_0(\mathbf{x}_0) + S_{EL}(\mathbf{x}_0; \mathbf{x}, t)) \quad (18)$$

as a min-plus path integral [12, 13] in a manner **analogous to Feynman's path integral**, but in the min-plus analysis, a non-linear analysis we have developed [14, 15] following Maslov [16, 17].

These statistical Hamilton-Jacobi equations (equations (14-17) of theorem 1) correspond to a set of classical particles, without interaction between each other and subjected to an external potential field $V(\mathbf{x}_G)$, and of which we only know, at the initial time, the probability density $\rho_0(\mathbf{x}_G)$ and the velocity field $\mathbf{v}_0(\mathbf{x}_G)$ through the intermediary of the initial action $S_0(\mathbf{x}_G)$ (with $\mathbf{v}_0(\mathbf{x}_G) = \frac{\nabla S_0(\mathbf{x}_G)}{m}$).

These are non-discerned prepared classical particles like the non-discerned prepared quantum particles. For these non-discerned prepared classical particles, the velocity of the center-of-mass of the classical particle is given at each point (\mathbf{x}_G, t) by:

$$\mathbf{v}(\mathbf{x}_G, t) = \frac{\nabla S(\mathbf{x}_G, t)}{m} \quad (19)$$

This equation shows that the solution $S(\mathbf{x}_G, t)$ to the Hamilton-Jacobi equations (14) defines the velocity field at any point (\mathbf{x}, t) from the velocity field $\frac{\nabla S_0(\mathbf{x})}{m}$ at the initial time. Thus, if given the initial position $X_G(0)$ of the center-of-mass of an prepared non-discerned classical particle, we deduce from (19) the trajectory $X_G(t)$ of the center-of-mass of the particle by the evolution equation:

$$\frac{dX_G(t)}{dt} = \frac{\nabla S(\mathbf{x}_G, t)}{m} \Big|_{\mathbf{x}_G=X_G(t)} \quad (20)$$

The Hamilton-Jacobi action $S(\mathbf{x}_G, t)$ is thus a field which **drives the motion of the center-of-mass of the classical particle**.

3.3 The dBB interpretation of the external wave function

To interpret the external wave function, we will use the mathematical continuity between the density and action of quantum mechanics and the density and action of classical mechanics, solutions to the Hamilton-Jacobi statistical equations. The **classical particles** which verify the Hamilton-Jacobi statistical equations have two properties:

- They are prepared non-discerned because we do not know at the initial time the position of their centers of mass, but only their initial distribution $\rho_0(\mathbf{x}_G)$. In classical mechanics, we remove the indeterminacy by adding the initial position of the center-of-mass $X_G(0)$.
- They are piloted by the gradient of the Hamilton-Jacobi action, which corresponds to a field that *pilots* the center-of-mass with the equation (20).

The mathematical continuity incites a continuity of interpretation by extending these two properties to the centers of mass of quantum particles that verify the Schrödinger equations of the external wave function:

- They are prepared non-discerned because we do not know at the initial time the position of their centers of mass, but only their initial distribution $\rho_0(\mathbf{x}_G)$. As in classical mechanics, we remove the indeterminacy by giving the initial position of the center-of-mass $X_G^h(0)$.
- The centers of mass of quantum systems are guided by the gradient of the quantum action, which corresponds to a field that “pilots” the center-of-mass by the equation:

$$\mathbf{v}^h(\mathbf{x}_G, t) = \frac{1}{m} \nabla S^h(\mathbf{x}_G, t) \quad (21)$$

Thus, the external wave function *pilots* the centers of mass of the quantum systems and corresponds to the **De Broglie-Bohm interpretation** [1, 6] restricted to the external wave function only. It is therefore necessary to add the initial position $X_G^h(0)$ of the center-of-mass as well as the equation that gives its evolution:

$$\frac{dX_G^h(t)}{dt} = \frac{1}{m} \nabla S^h(\mathbf{x}_G, t)|_{\mathbf{x}_G=X_G^h(t)} \quad (22)$$

When the potential $V(\mathbf{x}_G)$ can be considered as a linear gravitational potential, the Schrödinger equations (7-8) can be solved explicitly and one proves that the dBB trajectories converge to Newton trajectories [8, 9]. For our approach, the external wave function has the same meaning as the total wave function for the dBB theory : the wave function of the particle pilots its center-of-mass.

Louis de Broglie developed the **wave pilot theory** in 1926 for the total wave function. He presented it to the Solvey Congress in 1927 [18] as part of the theory of the double solution he was working on. However, he abandoned it a few years later. David Bohm rediscovered it in 1952, it is then also known as **Bohmian mechanics** [19]. It represents a causal, realistic and observer-free alternative to the Copenhagen interpretation [8, 20]. Louis de Broglie then resumed his work on this theory, which we call dBB theory. Since 2000, many studies have been carried out to develop and deepen the dBB approach both in theory and in applications.

Numerous simulations of dBB trajectories have been carried out by Sanz and Miret-Artès’ teams [21, 22] or for quantum chemistry applications [23]. The development of weak measurements paves the way for an possible experimental validation of these trajectories [24]. The approaches of Bohmian mechanics to be compatible to the quantum field theory are numerous [25, 26]. A synthesis of the dBB theory can be found in the books of Bricmont [27, 28] or of [29, 30] and in Goldstein’s article [19] in the Stanford Encyclopedia of Philosophy.

4 The internal wave function: the corpuscle of the field-corpuscle duality

For a quantum system corresponding to an atom or a molecule, the external wave function $\psi(\mathbf{x}_G, t)$ is defined in the laboratory reference frame while the relative wave function $\varphi(\mathbf{x}'_1, \mathbf{x}'_2, \dots, \mathbf{x}'_N, t)$ is defined in the center-of-mass reference frame.

Definition 2 - *From the relative wave function $\varphi(\mathbf{x}'_1, \mathbf{x}'_2, \dots, \mathbf{x}'_N, t)$ with $N > 1$ and the position of the center-of-mass $X_G^h(t)$, we define at each time, t , **the internal wave function** $\Phi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N, t)$ in the laboratory reference frame by the equation:*

$$\Phi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N, t) \equiv \varphi(\mathbf{x}_1 - X_G^h(t), \mathbf{x}_2 - X_G^h(t), \dots, \mathbf{x}_N - X_G^h(t), t) \quad (23)$$

We therefore make the following assumption :

Hypothesis 2 - Hypothesis of the double scale theory : The state of an N -body quantum system is simultaneously described by two wave functions in laboratory reference frame :

- the external wave function : $\psi(\mathbf{x}_G, t)$, a wave related to its center-of-mass : $X_G^h(t), t$
- the internal wave function : $\Phi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N, t)$.

Let us complete the definition of the internal wave function for a quantum system corresponding to a single particle, $N = 1$. This particle admits an external wave function $\psi(\mathbf{x}_G, t)$ with a center-of-mass $X_G^h(t)$ and a relative function $\varphi(\mathbf{x}', t)$. We define at each time, t , its **internal wave function** $\Phi(\mathbf{x}, t)$ in the laboratory reference frame by the equation:

$$\Phi(\mathbf{x}, t) \equiv \varphi(\mathbf{x} - X_G^h(t), t). \quad (24)$$

There are therefore two wave functions in the laboratory reference frame, the external wave function and the internal wave function. We have shown that the external wave function is a field that drives the center-of-mass of the corpuscle. Let us now study the convergence of the internal wave function when $\hbar \rightarrow 0$.

4.1 Convergence of the internal wave function

As the size of a single particle tends to zero with h , the density $|\Phi(\mathbf{x}, t)|^2$ of its internal wave function converges mathematically to the Dirac distribution $\delta(\mathbf{x} - X_G(t))$.

More generally, if this internal wave function corresponds to an atom or a molecule, its size depends on h and will tend towards zero if we make $h \rightarrow 0$. We deduce that the density $|\Phi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N, t)|^2$ of the internal wave function converges mathematically to the Dirac distribution at point $\mathbf{x} = X_G(t)$ when we make $h \rightarrow 0$. This convergence is completely different from the external wave function

These convergences are postulated and not demonstrated for the general case. However for particular cases, the demonstration is possible, as we recall below for the coherent states of the two dimensional harmonic oscillator. For coherent states, let us consider the time dependent Schrödinger equation with the potential $V(\mathbf{x}) = \frac{1}{2}m\omega^2(x^2 + y^2)$ and the initial wave function:

$$\Psi_0(\mathbf{x}) = (2\pi\sigma_h^2)^{-\frac{1}{2}} e^{-\frac{(\mathbf{x}-\mathbf{x}_0)^2}{4\sigma_h^2} + i\frac{m\mathbf{v}_0 \cdot \mathbf{x}}{\hbar}} = \sqrt{\rho_0^h(\mathbf{x})} \exp\left(i\frac{S_0^h(\mathbf{x})}{\hbar}\right) \quad (25)$$

where $\rho_0^h(\mathbf{x}) = (2\pi\sigma_h^2)^{-1} e^{-\frac{(\mathbf{x}-\mathbf{x}_0)^2}{2\sigma_h^2}}$ and $S_0(\mathbf{x}) = S_0^h(\mathbf{x}) = m\mathbf{v}_0 \cdot \mathbf{x}$ are respectively the initial density and action, with $\sigma_h = \sqrt{\frac{\hbar}{2m\omega}}$ and where \mathbf{x}_0 and \mathbf{v}_0 are the initial position and velocity of the center of the wave packet.

Therefore, the wave function $\Psi(\mathbf{x}, t)$ is:

$$\Psi(\mathbf{x}, t) = (2\pi\sigma_h^2)^{-\frac{1}{2}} e^{-\frac{(\mathbf{x}-\mathbf{x}(t))^2}{4\sigma_h^2} + i\frac{m\mathbf{v}(t) \cdot \mathbf{x} - g^h(t)}{\hbar}} \quad (26)$$

where $\mathbf{x}(t)$ and $\mathbf{v}(t)$ correspond to the position and velocity of a classical particle subjected to the potential $V(\mathbf{x}) = \frac{1}{2}m\omega^2(x^2 + y^2)$ where $g^h(t) = \int_0^t (\hbar\omega + \frac{1}{2}m\mathbf{v}^2(s) - \frac{1}{2}m\omega^2\mathbf{x}^2(s))ds$. We deduce the value of the density $\rho^h(\mathbf{x}, t)$ and of the action $S^h(\mathbf{x}, t)$:

$$\rho^h(\mathbf{x}, t) = (2\pi\sigma_h^2)^{-1} e^{-\frac{(\mathbf{x}-\mathbf{x}(t))^2}{2\sigma_h^2}} \quad \text{and} \quad S^h(\mathbf{x}, t) = m\mathbf{v}(t) \cdot \mathbf{x} - g^h(t). \quad (27)$$

We deduce the following theorem by considering the limit $\hbar \rightarrow 0$ of equation (27).

THEOREM 2 - Let the coherent states of the two-dimensional harmonic oscillator defined by equations (25). When \hbar tends to 0, then σ_h tends to 0, and the density $\rho^h(\mathbf{x}, t)$ and the action $S^h(\mathbf{x}, t)$, converge respectively to

$$\rho(\mathbf{x}, t) = \delta(\mathbf{x} - \mathbf{x}(t)) \quad \text{and} \quad S(\mathbf{x}, t) = m\mathbf{v}(t) \cdot \mathbf{x} - g(t) \quad (28)$$

with $g(t) = \int_0^t (\frac{1}{2}m\mathbf{v}^2(s) - \frac{1}{2}m\omega^2\mathbf{x}^2(s))ds$ and where $S(\mathbf{x}, t)$ is the Hamilton-Jacobi singular action.

The wave function therefore converges to the motion $\mathbf{x}(t)$ of a single classical **oscillator**. This classical particle is **discerned** because it is completely defined by its initial condition initial condition \mathbf{x}_0 and its initial action $S_0(\mathbf{x}) = m\mathbf{v}_0 \cdot \mathbf{x}$.

Note that one should speak of *the center-of-mass of the classical particle* rather than of *the classical particle*. In this framework, equation (28) corresponds to the center-of-mass equation and is written :

$$\rho(\mathbf{x}_G, t) = \delta(\mathbf{x}_G - X_G(t)) \quad \text{and} \quad S(\mathbf{x}_G, t) = mX'_G(t) \cdot \mathbf{x}_G - g(t) \quad (29)$$

with initial position $X_G(0) = \mathbf{x}_0$ and initial velocity $X'_G(0) = \mathbf{v}_0$.

4.2 Interpretation of the internal wave function for an elementary particle

The interpretation that we propose for the internal wave function is the one proposed by Schrödinger in 1926 [31] and at the Solvay congress of 1927 [32]. It is based on the solution of the coherent states of the harmonic oscillator, which is consistent with the second quantization and with the introduction of the creation and annihilation operators. These coherent states of the harmonic oscillator are very particular, as Schrödinger pointed out [31]: “*Our wave packet always remains grouped, and does not spread over an increasingly large space over time, as do, for example, wave packets that we are used to in optics.*” This wave packet corresponds to a single particle (soliton) which keeps its shape and whose center-of-mass follows a periodic trajectory identical to that of the center-of-mass of a classical harmonic oscillator. Schrödinger thinks that this interpretation extends to the electron of the hydrogen atom [31]: “*It is certain that it is possible to construct by a process quite similar to the previous one, wave packets gravitating on Kepler ellipses at a large number of quanta and forming the wave image of the electron of a hydrogen atom; but in this case the difficulties of calculation will be much greater than in the particularly simple example that we have treated here and which from this point of view is almost a classroom exercise.*”

Hypothesis 3 - Schrödinger conjecture for an elementary particle: *An elementary particle, such as a free electron or a bound electron in a hydrogen atom, can be considered as an extended particle whose density is given by the square $|\Phi(\mathbf{x}, t)|^2$ of the internal wave function in the laboratory reference frame.*

The position $X_G(t)$ of the center-of-mass of the particle is then obtained from the internal wave function by:

$$X_G(t) = \int \mathbf{x} |\Phi(\mathbf{x}, t)|^2 d\mathbf{x} \quad (30)$$

The simultaneous existence of the two wave functions makes quantum mechanics complete. Indeed, it is not necessary to add the initial position of the center-of-mass $X_G(0)$ as in the de Broglie-Bohm interpretation of the external wave function because it is defined by the initial internal wave function: $X_G(0) = \int \mathbf{x} |\Phi_0(\mathbf{x})|^2 d\mathbf{x}$.

The dBB theory can be seen as a specific case of the double-scale theory if the internal function is reduced to a point, the center-of-mass of the particle; i.e., $\Phi(\mathbf{x})$ is a Dirac function : $\Phi(\mathbf{x}, t) = \delta(\mathbf{x} - X_G(t), t)$.

Remark 1 *The change of variable $\Phi(\mathbf{x}, t) \equiv \varphi(\mathbf{x} - X_G^h(t), t)$ is very important. Indeed, if the average is compute with $\varphi(\mathbf{x}, t)$, then, by definition, we obtain :*

$$\int \mathbf{x} |\varphi(\mathbf{x}, t)|^2 d\mathbf{x} = 0 \quad (31)$$

Remark 2 - *The hypothesis of an extended particle obliges to introduce forces to maintain the cohesion of the particle. A model of the extended and deformable electron is proposed by Poincaré in his famous Palermo memoir [33]. Dirac makes the same argument in his article of 1962 “An extensible model of the electron” [34]. It is also the basis of Delmelt’s work on the size of the electron [35].*

The spatial extension of the electron is also consistent with the introduction of cut-offs in quantum electrodynamics (QED) to eliminate the infinities due to the hypothesis of point particles.

4.3 Interpretation of the internal wave function for a system of N particles

At the Solvay Congress in 1927, Schrödinger generalized his interpretation for the wave function of a single particle to that of a wave function of N particles in configuration space:

“I found the following way of looking at things useful; it may be a little naïve but it is easy to grasp. The classical system of material points does not really exist, but there is something that continuously fills all the space [...] the real system is a composite image of the classical system in all its possible states, obtained by using $\phi\phi^$ as a “weight function”. The systems to which the theory is applied are classically composed of a large number of charged material points. As we have just seen, the charge of each of these points is distributed continuously through space and each charge point e provides the contribution of the $e \int \phi\phi^* dx dy dz$ to the charge of the quarterly volume element $dx dy dz$. As $\phi\phi^*$ generally depends on time, these charges vary.”*

Schrödinger took up this interpretation in 1952 [36] and was strongly criticized by the Copenhagen school and in particular by Born in his 1954 Nobel lecture [37]:

“Schrödinger thought that his wave theory made it possible to return to deterministic classical physics. He proposed (and he has recently emphasized his proposal anew), to dispense with the particle representation entirely, and instead of speaking of electrons as particles, to consider them as a continuous density distribution $|\psi|^2$ (or electric density $e|\psi|^2$).”

The criticism related to the contradiction between Schrödinger’s definition of the particles as “narrow wave packets” and the fact that the external function spreads out in time. One can understand the criticism of such an interpretation if one does not differentiate between the internal and external wave functions; but this argument is no longer valid for the internal wave function.

We extend Schrödinger’s approach by postulating that the internal wave function of an N -body system is the product of N narrow wave packets each corresponding to a body of the system. We make the following assumption.

Hypothesis 4 - Generalized Schrödinger conjecture for a system of N particles - *To each internal wave function in the laboratory reference frame $\Phi(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots, \mathbf{x}_N, t)$ of a quantum system composed of N particles, one can associate N individual time-dependent wave functions in the laboratory reference frame $\Phi^j(\mathbf{x}, t)$ such that $|\Phi^j(\mathbf{x}, t)|^2$ represents the density of the particle $j = 1..N$. The second part of the Hypothesis is to assume the following equation:*

$$\Phi(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots, \mathbf{x}_N, t) = \prod_{j=1}^N \Phi^j(\mathbf{x}_j, t), \quad (32)$$

This last assumption consider that the extended particles, i.e. the individual wave functions $\Phi^i(\mathbf{x}, t)$, do not overlap in real space and therefore have disjoint supports : $\Phi^j(\mathbf{x}, t)\Phi^i(\mathbf{x}, t) = 0$ for all $i \neq j$. This is a property that is also verified by the individual relative Schrödinger wave functions $\varphi^j(\mathbf{x}, t) = \Phi^j(\mathbf{x} + X_G^h(t), t)$.

Remark 3 - *With this assumption, the internal wave function $\Phi(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots, \mathbf{x}_N, t)$ in the configuration space is replaced by N individual (internal) wave functions in the real three dimensional space.*

Remark 4 - *Hypothesis 4 is for example valid for the nuclei of the atoms of a molecule. This assumption is not respected for the stationary wave functions of the electrons of an atom because the orbitals can overlap. Nevertheless, we think that this assumption is valid if the electrons wave functions are time-dependent internal wave functions.*

We complete the hypothesis 4 with the following hypothesis:

Hypothesis 5 - We assume that the N individual time-dependent wave functions $\varphi^j(\mathbf{x}_j, t)$ are solutions to the N individual non linear Schrödinger equations:

$$i\hbar \frac{\partial \varphi^j(\mathbf{x}_j, t)}{\partial t} = -\frac{\hbar^2}{2m_j} \Delta_{\mathbf{x}_j} \varphi^j(\mathbf{x}_j, t) + \left(\sum_{\substack{i=1 \\ i \neq j}}^N \int d\mathbf{x}_i |\varphi^i(\mathbf{x}_i, t)|^2 U_{ji}(|\mathbf{x}_j - \mathbf{x}_i|) \right) \varphi^j(\mathbf{x}_j, t) \quad (33)$$

with the initial conditions:

$$\varphi^j(\mathbf{x}_j, 0) = \varphi_0^j(\mathbf{x}_j) = \Phi_0^j(\mathbf{x}_j + X_G^h(0)). \quad (34)$$

Because the size of each particle $i = 1 \dots N$ (i.e. the support of φ^i) is very small compared to the size of a atom or molecule, we can assume that we have for all t , $|\Phi^i(\mathbf{x}, t)|^2 \simeq \delta(\mathbf{x} - X^i(t))$ and $|\varphi^i(\mathbf{x}, t)|^2 \simeq \delta(\mathbf{x} - \mathbf{x}^i(t))$ where $\mathbf{x}^i(t) = \int \mathbf{x} |\varphi^i(\mathbf{x}, t)|^2 d\mathbf{x}$ is the position of the center-of-mass of particle i in the center-of-mass reference frame and $X^i(t) = \int \mathbf{x} |\Phi^i(\mathbf{x}, t)|^2 d\mathbf{x}$ in the laboratory reference frame with $X^i(t) = \mathbf{x}^i(t) + X_G^h(t)$. The integral of equation (33) can then be approximated by: $\int d\mathbf{x}_i |\varphi^i(\mathbf{x}_i, t)|^2 U_{ji}(|\mathbf{x}_j - \mathbf{x}_i|) \approx U_{ji}(|\mathbf{x}_j - \mathbf{x}^i(t)|)$ and we obtain the following system of equations:

$$i\hbar \frac{\partial \varphi^j(\mathbf{x}_j, t)}{\partial t} = -\frac{\hbar^2}{2m_j} \Delta_{\mathbf{x}_j} \varphi^j(\mathbf{x}_j, t) + \left(\sum_{\substack{i=1 \\ i \neq j}}^N U_{ji}(|\mathbf{x}_j - \mathbf{x}^i(t)|) \right) \varphi^j(\mathbf{x}_j, t) \quad (35)$$

$$\mathbf{x}^i(t) = \int \mathbf{x} |\varphi^i(\mathbf{x}, t)|^2 d\mathbf{x} \quad (36)$$

with the initial conditions :

$$\varphi^j(\mathbf{x}_j, 0) = \varphi_0^j(\mathbf{x}_j) \quad (37)$$

The approximate calculation of equations (35-37) can be obtained from Ehrenfest's theorem.

There are many arguments, both experimental and theoretical, in favor of this generalized Schrödinger interpretation for the internal wave function (Hypothesis 4):

- It is fundamentally compatible with the methodology of **molecular dynamics**. Indeed, if we separate the nuclei from the electrons in a molecule, the nuclei will have internal wave functions verifying the two Schrödinger conjectures.
- It is also fundamentally compatible with the first Hohenberg-Kohn theorem[38], which is the basis of **density functional theory**, which states that a given electron density corresponds to a unique wave function.
- It is also in agreement with the recent (2019) experiment by Mineev, Devoret et al. [39] on “*the jump from ground state to an excited state of a three-level superconducting artificial atom*”. This experiment seems to prove Schrödinger right in his 1952 discussion with the Copenhagen school on quantum jumps, given that “*the experimental results show that the evolution of each jump made is continuous, coherent and deterministic*”. As Devoret explains: “*Our experimental results show that quantum jumps are unpredictable and discrete (as Bohr thought) over long periods of time, they can be continuous (as suggested by Schrödinger) and predictable over short periods of time*”.

5 A new reading of quantum mechanics

This scale-dependent dual interpretation is a framework for reading quantum mechanics in a simple way and also explains the choices of other interpretations that did not distinguish between external and internal wave functions.

The wave-corpuscule duality corresponds to the simultaneous existence of two wave functions linked together, at each instant t , by the position of the center-of-mass $X_G^h(t)$: while the external wave

function corresponds to the wave (field) and the internal wave function corresponds to the corpuscle. **Theoretically**, we therefore have field AND corpuscle. **Experimentally**, a coherent source of particles is a set of particles which all have the same external wave function, but whose internal wave functions are different. Thus in the slit diffraction, double slit interference, Stern and Gerlach spin measurement, tunneling effect and EPR-B experiments, the preparation of the quantum system is represented by the external wave function alone. The internal wave function, which is unknown, explain the statistical results of Born's interpretation.

Quantum mechanics is complete if the state of a quantum system corresponds to the simultaneous existence of both wave functions, incomplete otherwise. The position of the center-of-mass in the external wave function that must be added in the de Broglie-Bohm interpretation is obtained from the internal wave function by the equations (30) and (36). We can thus interpret Everett's multiple worlds as the set of internal wave functions compatible with the external wave function.

The measurement corresponds to the impact of the internal wave on the detection screen. It is thus the internal wave function that is involved in the collapse of the wave packet. The Copenhagen interpretation, which does not differentiate between the external and internal waves, cannot therefore see that the reduction of the wave packet concerns mainly the internal wave function. For the same reason, the GRW interpretation [40] requires an objective non-linear perturbation. This perturbation exists because the particle is stopped by the detection screen following a non-linear absorption phenomenon. In the double slit experiment, **the external wave function passes through both slits while the internal wave function passes through only one slit.**

Heisenberg's inequalities correspond, for the external wave function, to **uncertainty relations** on the positions and velocities of the centers of mass of an ensemble of molecules admitting this same wave function. For the internal wave function, the Heisenberg equalities correspond to **indeterminacy relations** on the different positions and velocities of this extended particle [9].

The non-local hidden variable of Bell's theorem and the EPR-B experiment concerns the external wave function [9] while the position of the impacts is a **local measured variable** corresponding to the internal wave function.

This scale-dependent dual solution can be tested experimentally by an asymmetric double slit experiment such as those we proposed a few years ago [41] and are currently pursuing [42].

6 Conclusion

We have proposed an experimentally testable interpretation of non-relativistic quantum mechanics that gives a new understanding of the links between quantum and classical mechanics. This scale-dependent double solution seems strongly consistent with the specifications of Louis de Broglie who wrote in 1971:

"I introduced, under the name of "theory of the double solution", the idea that it was necessary to differentiate between two solutions, distinct but intimately connected to the wave equation, one of which I called the u-wave, being a real and non-normalizable physical wave with a local accident defining the particle and represented by a singularity, the other one, Schrödinger's ψ wave, normable and without singularity, which would be only a representation of probabilities." [3]

Our external wave function corresponds to the Schrödinger's ψ wave and our internal wave function corresponds to the u-wave.

We believe that this new reading grid allows us to extend this realistic interpretation to all quantum mechanics as well as to relativity. We are indeed preparing an extension of this double scale theory to the relativistic case by considering the Gordon decomposition of the Dirac equation as a convection current corresponding to the external wave function (large components) and a spin current corresponding to the internal wave function (small components) [43]. We will also propose a semi-classical gravity converging to the Newton gravity when $\hbar \rightarrow 0$ [44]. An extension to the second quantization [45] also seems possible by considering the external wave function as a field and the internal wave function as extended particles that can be created or annihilated by being bound to the excitations of this field [45].

This interpretation of quantum mechanics may also be considered as an answer to one of the Einstein's final texts on the interpretation of quantum mechanics (1953), *Elementary Considerations on the Interpretation of the Foundations of Quantum Mechanics* in homage to Max Born [46]:

“The first effort goes back to de Broglie and has been pursued further by Bohm with great perspicacity [...] The second attempt, which aims at achieving a "real description" of an individual system, based on the Schrödinger equation, has been made by Schrödinger himself. Briefly, his ideas are as follows. The ψ -function itself represents reality, and does not stand in need of the Born interpretation[...] From the previous considerations, it follows that the only acceptable interpretation of Schrödinger's equation up to now is the statistical interpretation given by Born. However, it does not give the "real description" of the individual system, but only statistical statements related to sets of systems.”

References

- [1] L. de Broglie, La mécanique ondulatoire et la structure atomique de la matière et du rayonnement, Journal de physique 8 (1927) 225–241, english translation in [18]. (1927).
- [2] L. de Broglie, Une tentative d'interprétation causale et non linéaire de la mécanique ondulatoire : La théorie de la double solution, Gauthier-Villars, 1956, english translation in [47]. (1956).
- [3] L. de Broglie, J. Andrade e Silva, La réinterprétation de la mécanique ondulatoire, Gauthier-Villars, 1971 (1971).
- [4] D. Fargues, Louis de Broglie's 'double solution', a promising but unfinished theory, Annales de la Fondation Louis de Broglie 42 (1) (2017) 9–18 (2017).
- [5] L. de Broglie, La Nouvelle Dynamique des Quanta, in: Electrons et Photons, Rapports du Cinquième Conseil de Physique Solvay, Gauthier-Villard, 1928, pp. 118–119, transled in [18] p.385 "The new dynamics of quanta" (1928).
- [6] D. Bohm, A Suggested interpretation of the quantum theory in terms of hidden variables., Phys.Rev. 85 (1952) 166–193 (1952).
- [7] D. Bohm, B. Hiley, The Undivided Universe, Routledge, 1993 (1993).
- [8] P. R. Holland, The Quantum Theory of Motion, Cambridge University Press, 1993 (1993).
- [9] M. Gondran, A. Gondran, C. Noûs, The two-scale interpretation: de broglie and schrödinger's external and internal wave functions, Annales de la Fondation Louis de Broglie 46 (1) (2021) 87–126 (2021).
- [10] E. Madelung, Quantentheorie in hydrodynamischer Form, Zeit. Phys. 40 (1926) 322–326 (1926).
- [11] F. Shimizu, K. Shimizu, H. Takuma, Double-slit interference with ultracold metastable neon atoms, Phys. Rev. A 46 (1992) R17–R20 (Jul 1992).
- [12] M. Gondran, A. Gondran, A synthetic interpretation: the double-preparation theory, Physica Scripta T163 (2014) 014029, hal-01109005 (Dec. 2014).
- [13] A. Kenoufi, M. Gondran, A. Gondran, Semi-Classical Limit and Least Action Principle Revisited with $(\min,+)$ Path Integral and Action-Particle Duality, Russian Journal of Mathematical Physics 27 (1) (2020) 61–75 (2020).
- [14] M. Gondran, Analyse MinPlus, C. R. Acad. Sci. Paris 323 (1996) 371–375 (1996).
- [15] M. Gondran, Schrödinger Equation and Minplus Complex Analysis, Russian Journal of Mathematical Physics 11 (2) (2004) 130–139 (2004).

- [16] V. Maslov, *Analyse Idempotente*, Edition Mir, 1989 (1989).
- [17] V.P. Maslov, S. Samborskii, *Idempotent Analysis*, *Advances in Soviet Mathematics* 13, American Mathematical Society, 1992 (1992).
- [18] G. Bacciagaluppi, A. Valentini, *Quantum Theory at the Crossroads: Reconsidering the 1927 Solvay Conference*, Cambridge University Press, 2009, arXiv:quant-ph/0609184 (2009).
- [19] S. Goldstein, *Bohmian Mechanics* — *Stanford Encyclopedia of Philosophy*, <https://plato.stanford.edu/entries/qm-bohm/>, [Online; accessed 12-2022] (2021).
- [20] S. Goldstein, *Quantum Theory without Observers—Part One*, *Physics Today* 51 (3) (1998) 42–46 (1998).
- [21] A. Sanz, F. Borondo, S. Miret-Artés, *Particle diffraction studied using quantum trajectories*, *J. Phys. C: Condens. Matter* 14 (2002) 6109–6145 (2002).
- [22] A. Sanz, S. Miret-Artés, *A Trajectory Description of Quantum Processes. I. Fundamentals, A Bohmian Perspective*, Springer, 2012 (2012).
- [23] R. Wyatt, *Quantum Dynamics with Trajectories: Introduction to Quantum Hydrodynamics*, Springer, 2005 (2005).
- [24] S. Kocsis, B. Braverman, S. Ravets, M. J. Stevens, R. P. Mirin, L. K. Shalm, A. M. Steinberg, *Observing the Average Trajectories of Single Photons in a Two-Slit Interferometer*, *Science* 332 (6034) (2011) 1170–1173 (2011).
- [25] W. Struyve, *Pilot-Wave Theory and Quantum Fields*, *Reports on Progress in Physics* 73 (10) (2010) 106001 (2010).
- [26] D. Dürr, S. Goldstein, R. Tumulka, N. Zanghi, *Bohmian Mechanics and Quantum Field Theory*, *Physical Review Letters* 93 (2004) 1–4 (2004).
- [27] J. Bricmont, *Making Sense of Quantum Mechanics*, Springer, 2016 (2016).
- [28] J. Bricmont, *Quantum Sense and Nonsense*, Springer, 2017 (2017).
- [29] T. Norsen, *Foundations of Quantum Mechanics*, Springer, 2017 (2017).
- [30] X. Oriols, J. Mompart, *Applied Bohmian Mechanics: From Nanoscale Systems to Cosmology*, Singapore: Pan Stanford Publishing, 2012 (2012).
- [31] E. Schrödinger, *Der stetige Übergang von der Mikro-zur Makromechanik*, *Naturwissenschaften* 14 (1926) 664–666 (1926).
- [32] E. Schrödinger, *La mécanique des ondes, Electrons et photons: rapports et discussions du cinquième Conseil de physique tenu à Bruxelles du 24 au 29 octobre 1927*. Institut international de physique Solvay (1928), Gauthier-Villars et cie., 1928 (1928).
- [33] H. Poincaré, *Sur la dynamique de l'électron*, *Rend. Circ. Matem. Palermo* 21 (1906) 129–175 (Dec. 1906).
- [34] P. Diarc, *An extensible model of the electron*, *Proc. R. Soc. Lond. A* 1332 (268) (1962) 57–47 (1962).
- [35] H. G. Dehmelt, *Experiments with an isolated subatomic particle at rest*, *Nobel Lecture* (1989) 1–13 (1989).
- [36] E. Schrödinger, *Are there quantum jumps?*, *Brit. J. Phil. Sc.* 322 (1952) 109–123; 233–242 (1952).

- [37] M. Born, Continuity, determinism, and reality, Kongelige Danske Videnskabernes Selskab, Matematiskfysiske Meddelelser 30 (2) (1955) 1–26 (1955).
- [38] P. Hohenberg, W. Kohn, Inhomogeneous Electron Gas, Phys. Rev. 136 (1964) B864–B871 (Nov 1964).
- [39] Z. K. Mineev, S. O. Mundhada, S. Shankar, P. Reinhold, R. Gutiérrez-Jáuregui, R. J. Schoelkopf, M. Mirrahimi, H. J. Carmichael, M. H. Devoret, To catch and reverse a quantum jump mid-flight, Nature 570 (2019) 200–204 (2019).
- [40] G. C. Ghirardi, A. Rimini, T. Weber, Unified dynamics for microscopic and macroscopic systems, Phys. Rev. D 34 (1986) 470–491 (Jul 1986).
- [41] M. Gondran, M. Božić, D. Arsenović, A. Gondran, Proposed Experiment with Rydberg Atoms to Study the Influence of Particle Size on Quantum Interference, Acta Physica Polonica A 112 (5) (2007) 805 (November 2007).
- [42] M. Gondran, A. Gondran, Variation of the double slit experiment to test the wave-particle duality, in progress (2023).
- [43] M. Gondran, A. Gondran, Double scale theory and relativity, in progress (2023).
- [44] M. Gondran, A. Gondran, Semiclassical gravity in the de broglie-bohm theory and the double scale theory, in submission, hal-03945789v1 (2022).
- [45] M. Gondran, A. Gondran, Double scale theory and second quantification, in progress (2023).
- [46] A. Einstein, Elementare Überlegungen zur Interpretation der Grundlagen der Quanten-Mechanik, Scientific Papers presented to Max Born, Olivier and Boyd; trad. fr. in A. Einstein, Œuvres choisies 1, Quanta, Seuil et CNRS, 1989. 368 (5) (1953) 409–547 (October 1953).
- [47] L. de Broglie, Non-linear wave mechanics. A causal interpretation, Elsevier Pub. Co., 1960 (1960).